

DESCRIPTION

TestDiet® PriLieve™ Bismuth Subsalsicylate Tablets are banana-flavored tablets with bismuth subsalsicylate, kaolin, pectin, and papain, used for the treatment of diarrhea and other gastrointestinal disorders in non-human primates.

Storage conditions are particularly critical to TestDiet® products, due to the absence of antioxidants or preservative agents. To provide maximum protection against possible changes during storage, store in a dry, cool location. Storage under refrigeration (2° C) is recommended. Maximum shelf life is six months. (If long term studies are involved, storing the diet at -20° C or colder may prolong shelf life.) Be certain to keep in air tight containers.

Product Forms Available*	Catalog #
1 gm Tablet (52.4 mg/tablet)	1812750
5 gm Tablet (262 mg/tablet)	1812749

*Other Forms Available By Request

INGREDIENTS

Active Ingredient: Bismuth Subsalsicylate (5.24%)
 Inactive Ingredients: Purified Primate Diet Base, Kaolin, Banana, Artificial Flavors, Silicon Dioxide, Pectin, Papain.

FEEDING DIRECTIONS

This tablet is intended as a supplement to a nutritionally complete diet and should only be fed under the direction of a veterinarian.

Actual results cannot be predicted and will vary by strain, husbandry, total diet, and general health of the animal.

Dosage: Speak to your veterinarian about proper dosing. Plenty of fresh, clean water should be available at all times.

This product is available with Enrofloxacin (Baytril): Bismuth Subsalsicylate + Enrofloxacin (Baytril) Tablets 5T4Z.

CAUTION:
 Perishable - store properly upon receipt.
 For laboratory animal use only; not for human consumption.

10/26/2007

NUTRITIONAL PROFILE¹

Protein, %		11.7	Minerals	
Arginine, %		0.45	Ash, %	4.3
Histidine, %		0.33	Calcium, %	0.38
Isoleucine, %		0.61	Phosphorus, %	0.32
Leucine, %		1.10	Phosphorus (available), %	0.32
Lysine, %		0.93	Potassium, %	0.55
Methionine, %		0.33	Magnesium, %	0.13
Cystine, %		0.25	Sulfur, %	0.09
Phenylalanine, %		0.61	Sodium, %	0.24
Tyrosine, %		0.65	Chlorine, %	0.40
Threonine, %		0.49	Fluorine, ppm	0.1
Tryptophan, %		0.14	Iron, ppm	129
Valine, %		0.73	Zinc, ppm	41
Alanine, %		0.35	Manganese, ppm	34
Aspartic Acid, %		0.82	Copper, ppm	7
Glutamic Acid, %		2.61	Cobalt, ppm	0.09
Glycine, %		0.25	Iodine, ppm	1.45
Proline, %		1.50	Chromium, ppm	1.83
Serine, %		0.71	Selenium, ppm	0.32
Taurine, %		0.00		
Fat (ether extract), %		2.4	Vitamins	
Fat (acid hydrolysis), %		2.4	Carotene, ppm	0.0
Cholesterol, ppm		0	Vitamin A, IU/g	7
Linoleic Acid, %		1.37	Vitamin D-3 (added), IU/g	7.0
Linolenic Acid, %		0.02	Vitamin E, IU/kg	35
Arachidonic Acid, %		0.00	Vitamin K (as menadione), ppm	0.3
Omega-3 Fatty Acids, %		0.02	Thiamin Hydrochloride, ppm	3
Total Saturated Fatty Acids, %		0.30	Riboflavin, ppm	3.5
Total Monounsaturated Fatty Acids, %		0.58	Niacin, ppm	35
Polyunsaturated Fatty Acids, %		1.39	Pantothenic Acid, ppm	10
			Folic Acid, ppm	7.0
			Pyridoxine, ppm	2.88
Fiber (max), %		4.1	Biotin, ppm	0.1
Neutral Detergent Fiber ² , %		3.2	Vitamin B-12, mcg/kg	7
Acid Detergent Fiber ³ , %		2.7	Choline Chloride, ppm	683
Nitrogen-Free Extract (by difference), %		67.6	Ascorbic Acid, ppm	3,363.4
Starch, %		3.31		
Glucose, %		0.00		
Fructose, %		0.00		
Sucrose, %		39.17		
Lactose, %		0.00		
Total Digestible Nutrients, %		43.4		
Energy (kcal/g)⁴		3.39		
From:	kcal	%		
Protein	0.468	13.8		
Fat (ether extract)	0.216	6.4		
Carbohydrates	2.702	79.8		

1. Based on the latest ingredient analysis information. Since nutrient composition of natural ingredients varies, analysis will differ accordingly. Nutrients expressed as percent of ration on an As-Fed basis except where otherwise indicated. Moisture content is assumed to be 10.0% for the purpose of calculations.
 2. NDF = approximately cellulose, hemicellulose and lignin.
 3. ADF = approximately cellulose and lignin.
 4. Energy (kcal/gm) - Sum of decimal fractions of protein, fat and carbohydrate x 4,9,4 kcal/gm respectively.

